

CRAB ARREST™

Domestic Granular

Pre-Emergent Dandelion and Large Crabgrass Weed Seed Germination Inhibitor with Corn Gluten

Apply Early Spring and Late Summer Before Weeds Appear

Guarantee: Corn Gluten meal 84.975%

Reg. No.: 32210 P.C.P. Act

Read the label before using

Keep out of reach of children

Caution Eye Irritant. Potential Sensitizer

20 kg

Lawn Life 935023 Airport Rd., Mono, ON Canada L9W 6C6
800-265-1605

CRAB ARREST may inhibit dandelion and large crabgrass seed germination when used in conjunction with a sound lawn maintenance program. Established dandelion and large crabgrass at the time of application will not be inhibited. This product is formulated for use in residential lawns where established perennial ryegrass and Kentucky blue grass is the predominant grass species.

DIRECTIONS FOR USE: May inhibit the germination of dandelions and large crabgrass seed in lawns where established perennial ryegrass and Kentucky bluegrass is predominant.

- Not recommended for use with a fertilizer containing nitrogen.
- Established dandelion and large crabgrass, at the time of application, will not be controlled. The inhibitory effect of the product to weed seeds generally dissipates in five weeks following application
- Do not apply under windy conditions
- Do not apply this product if over-seeding or re-sodding in the spring or fall.
- May inhibit weed seed germination when used in conjunction with a sound lawn maintenance program

BEFORE YOU APPLY: Determine size of area to be treated- measure length and width in metres or feet, then multiply to determine coverage area in square meters or feet (length X width). Next, subtract non-lawn areas.

- For best results, apply with a broadcast/rotary or drop fertilizer spreader at the recommended settings.
- Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to you lawn, and sweep any product that lands on your driveway, sidewalk or street back onto your lawn.
- **Apply at the rate of 11.25 kg per 100 sq. metres (23 lb per 1,000 sq. feet.)**

WHEN TO APPLY: Apply in spring before dandelion and large crabgrass seeds have germinated.

- For best results: Apply to established lawns twice a year; once in the early spring 2 weeks before weed seed germination, and once in the late summer or early fall after heat stress has passed.
- Apply when soil is moist and when rain is forecasted within 2 days of treatment. If rainfall does not occur within 2 days of treatment, irrigation is required. Excessive moisture at time of treatment may reduce the effectiveness of the product.

HOW TO APPLY: FILL & SET: Set the spreader to the appropriate setting indicated. Settings are figured at 11.25 kg per 100 sq. metres (23 lb per 1,000 sq. feet). Place your spreader on a driveway or other paved surface when filling. This guards against spillage on grass. Be sure the spreader is closed when filling it. **APPLY:** For even coverage, begin walking before you turn spreader on to the open position so as to avoid dropping product in one spot. To avoid gaps and streaks, spread around border of lawn first and then fill in centre with uniform, parallel passes. Walk at a steady pace for an even application. **NEVER APPLY BY HAND.** Water in application. Do not apply under windy conditions. Product application dates may vary for both the spring and late summer/early fall application from year to year according to weather conditions. The inhibitory effect of the product to weed seeds generally dissipates in five weeks following application. CRAB ARREST is to be applied to a mature lawn having a well-developed root mass. Do not apply the product on newly seeded grass as it may inhibit seeds from germinating, wait until after first mowing when root systems are established. If over-seeding or re-sodding in the spring, do not apply the product in the spring. If over-seeding or re-sodding in the fall, do not apply the product in the fall. For best results: Apply to established turf twice a year; once in the early spring and once in the late summer/early fall. Not recommended for use with a fertilizer containing nitrogen.

SPRING APPLICATION: Dandelion and Large Crabgrass Inhibition: Large crabgrass seeds germinate when soil temperature reaches 12.8 degrees C. Apply to established perennial ryegrass and Kentucky bluegrass turf in early spring two weeks before large crabgrass seed germination.

LATE SUMMER/EARLY FALL APPLICATION: Dandelion and Large Crabgrass Inhibition: Apply to established perennial ryegrass and Kentucky bluegrass turf in late summer or early fall after heat stress has passed.

APPLICATION RATE: Apply at a rate of 11.25 kg/100 metre square.

METHOD OF APPLICATION: 1. Measure the area to be applied by taking the length multiplied by the width of the total area in metres. One normal step is approximately one metre. Measure the size of any buildings or driveway in the same manner and subtract this amount from the total area. The remaining total will represent the grass area to be treated. 2. Apply CRAB ARREST using a rotary spreader for large areas. For smaller areas, apply CRAB ARREST using a small hand spreader. 3. Apply CRAB ARREST in a criss-cross pattern. Use one half the rate in each direction. Apply when soil is moist and when rain is forecasted within 2 days of treatment. If rainfall does not occur within 2 days of treatment, irrigation is required. Excessive moisture at time of treatment may reduce the effectiveness of the product.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. May cause sensitization. CRAB ARREST should not be applied if the applicator of member of the household has a sensitivity or allergy to corn. Avoid contact with skin, eyes or clothing. Avoid inhaling/breathing dusts. For good hygiene practice, wear a long-sleeved shirt, long pants, closed footwear and gloves when handling the product. In addition, it is recommended that a dust mask be worn when transferring the product to the spreader and during application.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Wash skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **IF INHALED:** Move to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferable by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE: Keep in cool, dry conditions, away from seed, fertilizer and other pesticides.

DISPOSAL: Do not re-use the empty container. Dispose in accordance with municipal or provincial regulations, if applicable. If no such regulations apply, wrap and dispose of empty container with household garbage.

IMPORTANT: Follow directions as indicated on this quality lawn care product as over application may cause injury to you lawn. Spreader application rates may vary due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owners' Manual.