



**Problem:** Anthracnose of Turf - *Colletotrichum graminicola* and *Microdochium bolleyi*



**Host Plants:** Primarily creeping bentgrass and annual bluegrass but may infect Kentucky bluegrass and tall fescue

**Description:** Anthracnose is a common disease of creeping bentgrass and annual bluegrass on golf course putting greens in Kansas. The disease is usually associated with turf suffering from heat injury in midsummer, but it can also develop in spring and fall. Anthracnose commonly occurs in conjunction with other diseases such as summer patch, Rhizoctonia brown patch and Pythium blight and is often a component in the so-called "bentgrass decline" complex.

Extensive damage may occur on bentgrass/annual bluegrass putting greens affected with anthracnose, and recovery from damage may be slow. Anthracnose may develop throughout the growing season, although it is more common in midsummer when cool-season turf is experiencing heat or drought stress. From a distance, affected bentgrass and/or annual bluegrass appears unthrifty and has a yellow or bronze cast. Affected turf wilts rapidly during midday and requires frequent irrigation. Anthracnose may initially be more severe on annual bluegrass. This results in a patchy appearance to the putting surface, as the bentgrass remains relatively unscathed while the annual bluegrass is killed. However, anthracnose may severely damage bentgrass greens with little or no annual bluegrass.

Individual plants affected with anthracnose first turn yellow and then die. Distinct leaf spots are not commonly formed by the anthracnose fungus. Instead, individual leaf blades fade from dark green to light green and then to yellow. There is no distinct region between healthy and diseased tissue as commonly observed with other leaf spotting fungi. The anthracnose fungus produces conspicuous black fruiting structures called acervuli on the leaves. The acervuli are abundant on dead tissue but may also form on green, apparently healthy leaves. The black hairy or spiny fruiting bodies are easily visible with a 10X hand lens.

The most severe damage occurs when the anthracnose fungus infects and colonizes the lower crown of the turfgrass plant. This is sometimes referred to as basal crown rot. Plants with crown rot are killed, resulting in a thinning of the turfgrass stand. Acervuli and small, black resting structures called sclerotia form on the decaying crown. The scattered structures are visible with a hand lens and appear as small pepper-like dots.

Anthracnose most often develops during the warm, humid weather of summer. Some pathologists consider the anthracnose fungus to be a weak pathogen at best, requiring substantial injury to the turfgrass plant for invasion. The presence of anthracnose often signifies significant injury to turf from environmental, chemical, or other pathological causes.

Two different anthracnose fungi are found in Kansas. *Colletotrichum graminicola* is the most common anthracnose fungus. This fungus has conspicuous spines or hairs that project up through the fruiting body. Spores of the fungus are crescent-shaped and may have one to two septations. A second anthracnose fungus called *Microdochium bolleyi* is sometimes associated with declining bentgrass. This fungus has spores similar to *Colletotrichum* but lacks the spiny protrusions of the fruiting structure.

**Recommendations:** Several cultural practices will help suppress anthracnose development. On greens with a history of the disease, consider a rigorous aerification program. Deep tine or core aerify in the fall. During the summer "open" the greens periodically by spiking, slitting, or hydro-jecting. Maintain a balanced fertilization

program. Do not starve the turf during the summer months. Very light fertilizations with a balanced fertilizer during the summer may help the turf withstand stresses and recover quickly from anthracnose damage. Low, frequent mowings enhance disease development. Consider raising mowing heights during periods of extreme turfgrass stress. Areas prone to localized dry spots are more susceptible to anthracnose damage. Follow a rigorous program to reduce damage from localized dry spots. Do not topdress during periods of turfgrass stress and anthracnose development. Topdressing damages plants and predisposes them to infection.

**References:**

1. Diseases of Turf, K-State Research and Extension Pub C-691
2. [Anthracnose of Bentgrass and Annual Bluegrass](#) K-State Research and Extension, Plant Pathology. Disease factsheet (9/96)

**Last Update:** 01/23/2004

---

[Extension Horticulture Home Page](#) | [Horticulture Information Center](#)