

AMBUSH[®] 500EC

Emulsifiable Concentrate Insecticide

AGRICULTURAL

GUARANTEE:

permethrin 500 g per litre

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

Store above 0 °C

REGISTRATION NO.: **14882** PEST CONTROL PRODUCTS ACT

NET CONTENTS: **1 Litre, 5 Litres**

AMVAC CHEMICAL CORPORATION
Suite 1200
4695 MacArthur Court
Newport Beach CA 92660
United States
323-264-3910

E93/E-5 706376/6 AMBUSH 500EC is a synthetic pyrethroid insecticide of low mammalian toxicity. It is a fast acting stomach and contact insecticide with no systemic or fumigant effect.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION: This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

Avoid splashing concentrate in eyes, on skin or on clothing. When applying this product in enclosed areas or with equipment capable of producing respirable spray droplets (less than 10 microns in size), an adequate respirator should be worn.

DO NOT APPLY BY GROUND EQUIPMENT WITHIN 15 M OF OR BY ANY AERIAL APPLICATION WITHIN 100M OF ANY BODY OF WATER, ESPECIALLY PRODUCTIVE FISHERIES WATER OR WATERFOWL HABITATS.

AERIAL APPLICATIONS SHOULD BE APPLIED IN LATE EVENING OR EARLY MORNING WHEN WIND SPEEDS ARE LESS THAN 10KPH. THIS PRODUCT IS VERY TOXIC TO FISH.

**DO NOT CONTAMINATE PONDS, LAKES, STREAMS OR RIVERS DURING SPRAYER FILLING OR RINSING OPERATIONS OR WHILE SPRAYING.
DO NOT APPLY WHEN WEATHER CONDITIONS FAVOUR DRIFT FROM THE TARGET AREA.**

Cover crop or crop treated with AMBUSH should not be used as green feed for animals.

This product is very toxic to bees; avoid spraying when bees are foraging. Spray deposit

should be dry before bees commence foraging in treated crop.

Re-entry to treated areas is permissible as soon as the spray deposit has dried.

This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal of waste or cleaning equipment.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in U.S., visit Croplife Canada's **website at : www.croplife.ca**.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

(1) Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.

(2) Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of container in accordance with provincial requirements. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING
CALL 1-323-264-3910**

AMBUSH[®] 500EC

Emulsifiable Concentrate Insecticide

AGRICULTURAL

GUARANTEE:

permethrin 500 g per litre

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

Store above 0°C

REGISTRATION NO.: **14882** PEST CONTROL PRODUCTS ACT

NET CONTENTS: 1 Litre, 5 Litres

AMVAC CHEMICAL CORPORATION

Suite 1200

4695 MacArthur Court

Newport Beach CA 92660

United States

323-264-3910

H01/E-6 AMBUSH 500EC is a synthetic pyrethroid insecticide of low mammalian toxicity. It is a fast acting stomach and contact insecticide with no systemic or fumigant effect.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION: This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

Avoid splashing concentrate in eyes, on skin or on clothing. When applying this product in enclosed areas or with equipment capable of producing respirable spray droplets (less than 10 microns in size), an adequate respirator should be worn.

DO NOT APPLY BY GROUND EQUIPMENT WITHIN 15 M OF OR BY ANY AERIAL APPLICATION WITHIN 100M OF ANY BODY OF WATER, ESPECIALLY PRODUCTIVE FISHERIES WATER OR WATERFOWL HABITATS.

AERIAL APPLICATIONS SHOULD BE APPLIED IN LATE EVENING OR EARLY MORNING WHEN WIND SPEEDS ARE LESS THAN 10KPH. THIS PRODUCT IS VERY TOXIC TO FISH.

**DO NOT CONTAMINATE PONDS, LAKES, STREAMS OR RIVERS DURING SPRAYER FILLING OR RINSING OPERATIONS OR WHILE SPRAYING.
DO NOT APPLY WHEN WEATHER CONDITIONS FAVOUR DRIFT FROM THE TARGET AREA.**

Cover crop or crop treated with AMBUSH should not be used as green feed for animals.

This product is very toxic to bees; avoid spraying when bees are foraging. Spray deposit

should be dry before bees commence foraging in treated crop.

Re-entry to treated areas is permissible as soon as the spray deposit has dried.

This product contains a petroleum distillate which is moderately to highly toxic to aquatic organisms. Avoid contamination of aquatic systems during application. Do not contaminate these systems through direct application, disposal of waste or cleaning equipment.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in U.S., visit Croplife Canada's **website at: www.croplife.ca**.

STORAGE

Store in a cool, dry, well ventilated area away from foodstuffs and out of the reach of children and animals. Keep product from freezing.

SPILL CLEANUP

Wear appropriate protective equipment (gloves, glasses, apron) when attempting to clean up the spill. If the container is leaking, secure leak and place the container into a drum or heavy gauge plastic bag. Contact Amvac Chemical Corporation at 1-323-24-3910 for further information.

For spills and leaks; contain the liquid with dikes of inert material (soil, clay, kitty litter etc.). Absorb the spill onto inert material and shovel into a sealable waste container. On hard surfaces - sprinkle spill area with detergent and scrub in a small quantity of water with a coarse broom. Let stand 10 minutes then absorb onto an inert material and shovel into the waste container. On soil - remove the top 15 cm of soil in the spill area and replace with fresh soil. Dispose of all waste including scrub brush in accordance with provincial requirements. For more information on the disposal of waste and the clean up of spills, contact the Provincial Regulatory Agency or the manufacturer.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

(1) Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.

(2) Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of container in accordance with provincial requirements. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING
CALL 1-323-264-3910**

DIRECTIONS FOR USE

Ground Equipment Only (For Aerial Application See Page 9)

PRECAUTION: Do not apply by ground equipment within 15m of any body of water especially within productive fisheries water or waterfowl habitats.

TOBACCO (GREENHOUSE PLANTS): Mix 8.25 mL in 50 L of water and apply 15 L of spray mixture to 100 m² of plant bed.

TOBACCO (FIELD): For control of **dark-sided cutworm** and **white cutworm** by methods given below, apply 140 mL/ha in 225 to 450 L of water using nozzle pressure of 175 to 350 kPa. Application of AMBUSH should be timed as close as possible to insect feeding activity which is greatest during the late evening and night.

COVER CROP TREATMENT: Apply once to rye or wheat cover when crop is 10 to 15 cm high, 4-5 days before ploughdown. Application should also be made to fence rows and a 15 m strip into nearby cover crop.

SOIL TREATMENT: Apply once to the soil 5 days before transplanting. (Under dry soil conditions, use 275 mL per hectare). Do not incorporate. Do not disturb the soil surface for at least 5 days following treatment since mixing of AMBUSH with soil will reduce its effectiveness. Application should also be made to fence rows and a 15 m strip into nearby cover crop.

POST-PLANTING TREATMENT: Apply once to transplants immediately prior to anticipated cutworm feeding damage. Under conditions of severe insect pressure, application should also be made to fence rows and a 15 m strip into nearby cover crops. A follow-up treatment may be necessary if there are late developing cutworms. (Do not apply less than 55 days before harvest).

Tank Mixing with DEVRINOL[®] 50-DF: if timing of AMBUSH 500EC application coincides with that of DEVRINOL[®] 50-DF for weed control, a tank mix may be used. Follow directions on AMBUSH 500EC and DEVRINOL[®] 50-DF labels for appropriate timing, rates and precautions.

CUTWORMS: (Army, Black, Dark-sided, Pale western, Red-backed and White): To control these insects attacking seedlings (up to 5 leaves) and/or transplants of **asparagus, beets, carrots, cereals, cole crops, corn, flax, lentils, lettuce, onions, peas, peppers, potatoes, rape, sugar beets and sunflowers** apply 140 mL/ha in sufficient water for good coverage. Under dry soil conditions or on muck soil, apply 225 to 300 mL/ha. Where cutworms are large (near full maturity - 2.5 to 4 cm), use 225 to 300 mL/ha. Applications should be made under warm, moist conditions in the evening or at night when cutworm activity is highest. **DO NOT DISTURB THE SOIL SURFACE FOR 5 DAYS AFTER TREATMENT.**

NOTE: AMBUSH 500EC will only control surface feeding or climbing stages of cutworms.

ASPARAGUS: Cutworms: To control cutworms during the cutting season, make 1 broadcast application to bare soil after cutting when damage is first noticed. Apply 140 mL/ha in sufficient water for good coverage. Apply in the evening or at night when cutworm activity is highest. Do not apply within 2 days of harvest.

CANOLA (RAPE): Crucifer Flea Beetle: Apply 70 to 140mL /ha.

Ground Application: Apply in sufficient water for good coverage when insects are present. Application should be made when the beetles are actively feeding. For severe infestations, use 140 mL per hectare.

Aerial Application: Application by air is permitted, provided there is no hazard of drift to other crops or to areas occupied by people or livestock. Apply specified rate in 20 -45 L water/ha. Can only be applied by air once per season.

COLE CROPS (Cabbage, Cauliflower, Broccoli, Brussels Sprouts): Cabbage Looper, Imported Cabbage Worm, Diamondback Moth Larvae and Crucifer Flea Beetle: Apply 70 to 140 mL in 400 to 675 L of water per hectare on a 7 to 10 day schedule when insects or insect

damage first appear. Add AGRAL[®] 90 wetting agent at 300 mL per 1000 L of water to improve wetting and coverage. For severe infestation use 140 mL per hectare. May be applied up to 3 days before harvest of cabbage, cauliflower and brussels sprouts and up to 7 days before harvest of broccoli.

CHINESE BROCCOLI: Cabbage Looper, Diamondback Moth Larvae, Imported Cabbage Worm: Apply 70 to 140 mL in 400 to 675 L of water per hectare on a 7 to 10 day schedule when insects or insect damage first appear. Add AGRAL[®] 90 wetting agent at 300 mL per 1000

L of water to improve wetting and coverage. For severe infestation use 140 mL per hectare. May be applied up to 7 days before harvest. Do not make more than 4 applications per year.

MUSHROOMS (Mushroom House and Adjacent Premise Areas): Mushroom flies (Sciarid and Phorid Adults) - Apply a fogging or aerosol treatment at the rate of 50 to 65 mL per 1 litre of water per 1,000 m³.

Preparation of the building prior to fogging:

- 1) Close all doors, windows and ventilators.
- 2) Lock or barricade all entrances, turn off pilot lights, post warning signs and take precautions to prevent persons and animals from entering the area.

Use prior to filling, during cool down, during spawning, up to pinning and between breaks.

NOTE:

Do not apply more than two applications between each break with maximum of 10 applications during cropping.

Do not apply within one day of picking.

Do not use when air temperature is less than 12°C or greater than 30°C. Use above 30°C could reduce the effectiveness of AMBUSH 500EC.

Depending upon fly populations and crop stage, treat once daily or as needed. However, do not make more than 20 applications prior to pinning of first break. Routine procedures for reducing background populations need to be carefully followed to ensure adequate adult fly control in the crop and to reduce the necessity for pesticide applications.

Consult provincial mushroom specialist for details of Integrated Pest Management programs to be followed for sciarid and phorid fly control. Length of exposure time should be limited to one hour, then ventilate the mushroom house. Use fans to ventilate in houses that do not have forced air circulation.

PAK-CHOI and CHINESE CABBAGE: Cabbage Looper, Imported Cabbage Worm, Diamondback Moth Larvae: Apply 140 mL in 400 to 675 L of water per hectare on a 7 to 10 day schedule when insects or insect damage first appear. Add AGRAL[®] 90 wetting agent at 300 mL per 1000 L of water to improve wetting and coverage. May be applied up to 3 days before harvest. Do not make more than 4 applications.

PEANUTS: Potato Leafhopper: Apply 140 mL/ha in sufficient water for thorough coverage when infestation exceeds 2 leafhoppers per plant. Repeat as required. Do not apply within 14 days of harvest.

PEPPERS: European Corn Borer: Apply 140 mL/ha when corn borer appears in your area and repeat every 7 days. Corn borer will be present from the end of July to mid-September. Do not apply within 1 day of harvest.

POTATOES: Colorado Potato Beetle, Potato Flea Beetle, Potato Leafhopper, Tarnished Plant Bug: Apply 140 to 200 mL/ha in sufficient water to give thorough coverage of plants. Use the higher rate for heavy infestations. Repeat as required.

Variegated (Climbing) Cutworm: Apply 140 mL per hectare when insects or damage appear - usually late July or during August, depending on location. Good control is dependent on spray penetration of dense foliage. Do not apply within 1 day of harvest.

European Corn Borer: Apply 140 mL/ha. Ground Application: Apply in sufficient water for good coverage. Apply when egg masses begin to hatch. Consult your crop specialists for appropriate spray schedule. Aerial Application: Application by air is permitted, provided there is no hazard of drift to other crops or to areas occupied by people or livestock. Apply specified rate in 20-45 L water/ha. Can only be applied by air once per season.

RADISH: Crucifer Flea Beetle: Apply 140 mL/ha when insects or insect damage appears. Repeat if required but do not apply within 2 days of harvest.

HORSERADISH: Crucifer Flea Beetle: Apply 140 mL/ha in sufficient water for good coverage at first sign of infestation. Repeat if required but do not apply within 2 days of harvest.

SNAP BEANS: European Corn Borer: Apply 200 mL/ha at the first sign of infestation. Repeat if necessary. Do not apply within 7 days of harvest.

SWEET CORN: European Corn Borer and Corn Earworm: Apply 200 to 275 mL in 325 to 450 L of water per hectare using a boom and nozzle arrangement to ensure thorough spray

coverage. Use the higher rate when severe insect pressure is anticipated. Spray not later than when first feeding is seen on foliage. For second brood borers in late plantings, apply before tassels show. For control of Corn Earworm, direct spray to ensure coverage of ears and silks. For the number and timing of repeat applications follow customary spray schedule in your region. Consult provincial spray calendar. **Fall Armyworm:** Apply 140 mL/ha. This treatment will not prevent internal cob damage if insect has penetrated ear. Do not apply within one day of harvest.

TOMATOES: Colorado Potato Beetle, Potato Flea Beetle, Tomato Hornworm and Variegated (Climbing) Cutworm - Apply 140 to 200 mL/ha in sufficient water to give thorough coverage. **Greenhouse Whitefly** - Apply 280 mL/ha. **Tomato Fruitworm (corn earworm)** - Apply 200 to 280 mL/ha using the higher rate for heavy infestations. **NOTE:** Tomato Fruitworm that has entered the fruit will not be controlled. Repeat above sprays as needed. Do not apply within 1 day of harvest.

TURNIPS: Crucifer Flea Beetle: Apply 140 mL/ha at the 2-8 leaf stage when insects or insect damage appears. Repeat if required but not later than the 8 leaf stage.

FRUITS: Apples, Pears, Peaches, Nectarines, Grapes and Blueberries: For all crops listed below, use sufficient water for thorough coverage. Follow provincial spray calendars for optimum timing of spray applications. Where a rate range is suggested, use the higher rate only when heavy insect pressure is anticipated.

NOTE: REPEATED APPLICATIONS ARE NOT ADVISED IN ORCHARDS WHERE INTEGRATED PEST MANAGEMENT IS IN EFFECT BECAUSE SEVERE REDUCTIONS OF BENEFICIAL ARTHROPODS MAY RESULT.

APPLES: Winter Moth, Eastern Tent Caterpillar - Apply 200 mL/ha. **Eye-spotted Budmoth:**

Apply 200 to 400 mL/ha. **Plum Curculio, White Apple Leafhopper** - Apply 275 to 400 mL/ha. **Tentiform Leafminer, Green Fruitworm, Apple Maggot, Leafrollers, Tarnished Plant Bug, Mullein Plant Bug, Lesser Appleworm, Codling Moth:** Apply 400 mL/ha. Follow provincial spray calendars for optimum timing of spray applications. Do not apply less than 7 days before harvest.

Apple Bark Borer, Dogwood Bark Borer - Mix 17 mL in 100 L of water with 2 L of Superior Oil. When monitoring indicates adults are active (late June to early August), apply to apple trunks as drench spray to runoff using a knapsack, handgun or airblast sprayer. A second application may be applied 14 days later if adults continue to be present. Do not apply within 7 days of harvesting.

PEARS: Eastern Canada - Pear Psylla (adults and nymphs), Codling Moth, Green Fruitworm: Apply 400 mL/ha. Do not apply less than 7 days before harvest. **British Columbia - Pre-bloom Pear Psylla (overwintering adults):** Apply 400 mL/ha up to 1 cm green stage of pear foliage growth. Do not use in British Columbia for control of nymphal stages of pear psylla. **Post-Bloom - Pear Psylla:** Apply 400 to 850 mL/ha. Do not apply less than 7 days before harvest at the 400 mL rate and less than 14 days before harvest at the 850 mL rate. **Codling Moth:** Apply 400 mL/ha. Do not apply less than 7 days before harvest.

PEACHES and NECTARINES: Oriental Fruit Moth: Apply 200 to 400 mL/ha. **Plum Curculio and Plant Bugs:** Apply 400 mL/ha. Do not apply less than 7 days before harvest.

PLUMS: Plum Curculio: Apply 425 mL/ha. Do not apply within 7 days of harvest.

GRAPES: Grape Berry Moth: Apply 275 mL/ha. **Grape Leafhopper:** Apply 140 mL/ha. Do not apply less than 7 days before harvest.

BLUEBERRIES (Vegetative year only): Thrips: Apply 140 mL/ha in one application between mid May and early June when plants are 1 to 2 cm tall.

ORNAMENTAL TREES AND SHRUBS AND COMMERCIAL NURSERIES: For insect control on trees and shrubs of ornamental value only (excluding Woodlands). **Balsam Fir, Douglas-Fir, Larch (Tamarack), Pine Trees and Spruce: Douglas-Fir Tussock Moth, White-Marked Tussock Moth, Spruce Budworm, Coneworm and Open Feeding Sawflies:** Apply 35-70 mL per hectare using application equipment available and dilutions suitable to equipment. Make one application per season. Applications should be calculated and made on the basis of volume per tree to ensure proper coverage while limiting off-target spray.

Other Coniferous Trees and Shrubs: Control of the above listed insects on coniferous tree and shrub species not listed above is acceptable provided the species is tested for phytotoxic effect prior to use. Spray a small, inconspicuous portion of the tree or shrub under test and wait for 24 hours to see if any adverse effects appear. **Deciduous Trees and Shrubs: Eastern Tent Caterpillar, Euonymus Webworm, Fall Webworm, Forest Tent Caterpillar and Open Feeding Aphids:** Apply 70 mL per 1000 L of water as a dilute spray (for mist blowers use 75 L of water). Apply when insects are first noticed.

Seedling Tree Nurseries: For cutworm control, apply 140 mL/ha. Application should be made under warm, moist conditions when larvae are small (up to 2.5 cm).

APPLICATION EQUIPMENT: The control achieved in any spray operation depends on the care taken during application. The correct type of equipment for the job should be selected. Powered, high-pressure sprayers and mist blowers are most suitable for treating large trees (or stands of trees) that are readily accessible from the ground.

High-pressure sprayers deliver a large volume of dilute spray mixture, and a sufficient quantity should be used to wet all of the foliage. Mist blowers are designed to apply a low volume of a concentrated insecticide mixture, and only enough should be used to moisten the foliage. Hand-pumped sprayers are usually adequate for treating medium-sized ornamentals up to 4.5 m in height. Mix AMBUSH 500 EC as for dilute spraying.

FORESTRY NURSERY GREENHOUSES: Conifer seedlings (spruce, pine, cedar, yew, juniper): Strawberry Root Weevil: Apply 50 mL of product per 100 L of water to runoff when adults emerge and before laying eggs. One application correctly timed per season should give control of adults. A second treatment may be made 21 days after monitoring shows continued presence of adults.

Willow and Poplar Trees: For the control of satin moth caterpillars (*Leucoma salicis*). Apply 70 mL Ambush 500 EC per 1000 L of water as a dilute spray. Apply as a ground application when insects are first noticed (late May to mid August) in a high pressure tree sprayer/mist blower. Application can be made twice per year to shade and ornamental trees in parks, school yards and residential areas.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use this product in a way that is consistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

NATURE OF RESTRICTION: To be used only in the manner authorized; contact provincial regulatory authorities about use permits which may be required.

RESTRICTED USES:

COMMERCIAL WOODLAND USE: Apply by ground application only. Use in Christmas tree plantations, high valued stands used for seed production and other high valued woodland areas that are extensively managed. Contact provincial or federal forestry and/or fisheries officials for advice before local spraying of woodlands. This product must be used in Woodlands under supervision of Provincial or Federal forestry officials.

Balsam Fir, Douglas-Fir, Larch (Tamarack), Pine Trees and Spruce: Douglas-Fir Tussock Moth, White-Marked Tussock Moth, Spruce Budworm, Spruce Coneworm and Open Feeding Sawflies: Apply 35-70 mL per hectare using application equipment available and dilutions suitable to equipment. Make one application per season. Applications should be calculated and made on the basis of volume per tree to ensure proper coverage while limiting off-target spray. **Spruce: Spruce Budmoth.** Apply 140 mL AMBUSH 500EC per hectare at beginning of hatch (mid May) using application equipment available and dilutions suitable to equipment. Precise application timing is important. Contact forestry officials for advice. Make one application per season. Applications should be calculated and made on the basis of volume per tree to ensure proper coverage while limiting off-target spray.

Balsam Fir: Hemlock Looper: Apply 35-70 mL AMBUSH 500EC per hectare between first and third instar using application equipment available and dilutions suitable to equipment. Make one application per season. Applications should be calculated and made on the basis of volume per tree to ensure proper coverage while limiting off-target spray.

Other Coniferous Trees and Shrubs: Control of the above listed insects on coniferous tree and shrub species not listed above is acceptable provided the species is tested for phytotoxic effect prior to use. Spray a small, inconspicuous portion of the tree or shrub under test and wait for 24 hours to see if any adverse effects appear.

Deciduous Trees and Shrubs: Eastern Tent Caterpillar, Euonymus Webworm, Fall Webworm, Forest Tent Caterpillar, and Open Feeding Aphids: Apply at the rate of 70 mL per 1000 L of water as a dilute spray (for mist blowers use 75 L of water). Apply when insects are first noticed. **Gypsy Moth.** Apply 35-70 mL AMBUSH 500EC per hectare between first and third instar using application equipment available and dilutions suitable to equipment. Make one application per season. Applications should be calculated and made on the basis of volume per tree to ensure proper coverage while limiting off-target spray.

Seedling Trees in Reforestation Areas: For Cutworm control, apply 140 mL/ha. Application should be made under warm, moist conditions and when larvae are small (up to 2.5 cm).

APPLICATION EQUIPMENT: The control achieved in any spray operation depends on the care taken during application. The correct type of equipment for the job should be selected. Powered, high-pressure sprayers and mist blowers are most suitable for treating large trees (or stands of trees) that are readily accessible from the ground. High-pressure sprayers deliver a large volume of dilute spray mixture, and a sufficient quantity should be used to wet all of the foliage. Mist blowers are designed to apply a low volume of a concentrated insecticide mixture, and only enough should be used to moisten the foliage. Hand-pumped sprayers are usually

adequate for treating medium-sized ornamentals up to 4.5 m in height. Mix AMBUSH 500 EC as for dilute spraying.

PRECAUTION: WHEN USING AS DIRECTED IN WOODLANDS, DO NOT APPLY WITHIN 100 M OF PRODUCTIVE FISHERY WATERS OR WATERFOWL HABITATS. SEE ADDITIONAL PRECAUTIONS ON THIS LABEL.

CLUSTER FLIES: Cluster flies are members of the fly family (Diptera) and resemble house flies. They are distinguished by their sluggish and erratic behaviour and by wings that overlap when at rest. They derive their name from their habit of entering a building in the fall and clustering together to overwinter. The search for protected places to overwinter begins as days shorten in mid-August. Warm sunny walls draw the flies as the climate cools during this search. This offers an opportunity to control the fly population outside the building.

Outdoor Control: Cluster flies: Prepare a spray mixture of 100 mL in 10 L of water. Spray this mixture to the outside walls, including under the eaves, soffits and window trim where applicable, of barracks, mess halls, drill halls, hangers, warehouses, equipment buildings, farm buildings and other buildings where cluster flies have become a nuisance. Apply at the approximate rate of 10 L of spray mixture per 100 m². The effective time of application is between mid-August and the end of September depending on weather conditions.

SPIDERS: Mix 100 mL in 10L of water and spray this mixture to areas where spiders congregate especially under eaves and soffits of buildings. Apply when heavy infestation of spiders are observed.

AERIAL APPLICATION

Generic Aerial Application Label Instructions - Directions for Use

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides. Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before using this product.

For aerial applications, ensure the aircraft is equipped and calibrated to deliver a uniform spray coverage with a minimum potential for drift. To ensure uniform application use an appropriate marking device. Apply in weather conditions that will not promote drift. Suggested conditions for good aerial application are moderate temperatures (less than 25 °C), moderate relative humidity (greater than 40%), and light winds (3 to 9 kph). Aerial drift is increased under certain meteorological conditions. Do not apply during periods of dead calm, when winds are gusty or when temperature inversion is likely (e.g. evening when warm air is rising from the crop or morning when sunshine warms the soil and air rises from the field). Do not apply by air in winds greater than 16 kph at flying height at the site of application.

Use only medium or coarse nozzles rated as delivering droplets of volume median diameter of 215 microns or greater. Examples: 6506 flat fan, CP 0.125 deflector nozzle with a low shear deflector angle (30 degrees), D12-56 disc-core.

AMBUSH may be applied twice per season by this method on snap beans. AMBUSH 500EC may be applied only once per season by this method for pale western cutworm and on potatoes and sweet corn.

PRECAUTION: DO NOT APPLY ANY AERIAL APPLICATION WITHIN 100 M OF ANY BODY OF WATER, ESPECIALLY PRODUCTIVE FISHERIES WATER OR WATERFOWL HABITATS. APPLY LATE EVENING OR EARLY MORNING WHEN WIND SPEEDS ARE LESS THAN 10 KPH.

POTATOES: Potato Flea Beetle, Colorado Potato Beetle, Potato Leafhopper, Tarnished Plant Bug: Apply 140 to 200 mL/ha in 20 to 45 L of water. **European Corn Borer:** Apply 140 mL/ha in 20 to 45 L of water.

PALE WESTERN CUTWORM: To control this insect attacking seedlings (up to 5 leaves) of cereals, corn, flax, potatoes, rape (canola), sugar beets, sunflowers, peas, lentils and other crops meeting the criteria of seedlings up to 5 leaves, apply 140 mL/ha in sufficient water for good coverage. Under dry soil conditions or where cutworms are large (near full maturity - 2.5 to 4 cm) use 225 to 300 mL per hectare. **NOTE:** AMBUSH 500 EC will only control surface feeding stages of cutworms.

SNAP BEANS: European Corn Borer: Apply 200 mL/ha in 20 to 45 L of water at the first sign of infestation. Do not apply within 7 days of harvest.

SWEET CORN: European Corn Borer and Corn Earworm: Apply 200 to 275 mL in 20 to 45 L of water per hectare. Use the higher rate when severe insect pressure is anticipated. Spray not later than when first feeding is seen on foliage. For second brood borers in late plantings, apply before tassels show. For control of Corn Earworm, spray to ensure coverage of ears and silks. **Fall Armyworm:** Apply 140 mL/ha. This treatment will not prevent internal cob damage if insect has penetrated ear. Do not apply within one day of harvest.

This product is very toxic to fish and aquatic organisms. For the protection of non-target habitats, overspray or drift to sensitive habitats must be avoided. A buffer zone of 100 metres is required between the downwind edge of the boom and the closest edge of sensitive aquatic habitats such as sloughs, ponds, prairie potholes, lakes, rivers, streams, and wetlands, (and wildlife habitat at the edge of these bodies of water). Do not contaminate these habitats when cleaning and rinsing spray equipment or containers.

Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

Clean and decontaminate protective clothing and application equipment regularly.

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for this product for the uses described on this Label were developed by persons other than Amvac Chemical Corporation and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Amvac Chemical Corporation itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used on the crop(s) listed on the above Label.

Accordingly, the User assumes all liability arising, and agrees to hold Amvac Chemical Corporation harmless from any claims based on efficacy and/or phytotoxicity in

connection with the use(s) described on this Label.

North American Ginseng - Fourlined Plant Bug: Apply 140-200 mL/ha of AMBUSH 500 EC in sufficient water. Maximum of two applications per year; ground only. Preharvest interval is 40 days. Application timing determined by scouting.

WASABI-Cabbage Looper, Imported Cabbage Looper, Diamondback Moth Larvae and Crucifer Flea Beetle: Ground Equipment only. Apply 70 to 140 mL in 400 to 675 L of water per hectare on a 7 to 10 day schedule when insects or insect damage first appear. Add AGRAL® 90 wetting agent at 300 mL per 1000 L of water to improve wetting and coverage. Do not apply within 7 days of harvest. Do not make more than 4 applications per year.

Resistance-Management Recommendations

For resistance management, please note that AMBUSH 500EC Insecticide contains a Group 3 insecticide. Any insect population may contain individuals naturally resistant to AMBUSH 500EC Insecticide and other Group 3 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay insecticide resistance:

Where possible, rotate the use of AMBUSH 500EC Insecticide or other Group 3 insecticides with different groups that control the same pests in a field.

Avoid application of more than the maximum number listed in the label and consecutive sprays of AMBUSH 500EC Insecticide or other insecticides in the same group in a season.

Use tank mixtures with insecticides from a different group when such use is permitted.

Insecticide use should be based on an IPM program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.

Monitor treated pest populations for resistance development.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance, contact company representatives at 1-323-264-3910.

AMBUSH is a registered trademark of Amvac Chemical Corporation.
Agral 90 and DEVRINOL are registered trademarks of Syngenta Crop Protection Canada, Inc.

Canadian Patent No. 1008455

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.